



UNIVERSITI PUTRA MALAYSIA

**DEVELOPING A DIET OF CATERPILLARS AND MEALWORMS
FOR MASS REARING OF THE PREDATORY BUG
CANTHECONIDEA FURCELLATA (WOLFF.)
(HEMIPTERA: PENTATOMIDAE)**

EVAWATY SRI ULINA

FP 2000 18

**DEVELOPING A DIET OF CATERpillARS AND MEALWORMS
FOR MASS REARING OF THE PREDATORY BUG
CANTHECONIDEA FURCELLATA (WOLFF.)
(HEMIPTERA: PENTATOMIDAE)**

EVAWATY SRI ULINA

**MASTER OF AGRICULTURAL SCIENCE
UNIVERSITI PUTRA MALAYSIA**

2000



**DEVELOPING A DIET OF CATERpillARS AND MEALWORMS
FOR MASS REARING OF THE PREDATORY BUG
CANTHECONIDEA FURCELLATA (WOLFF.)
(HEMIPTERA: PENTATOMIDAE)**

By

EVAWATY SRI ULINA

**Thesis Submitted in Fulfilment of the Requirements for the
Degree of Master of Agricultural Science in
the Faculty of Agriculture
Universiti Putra Malaysia**

September 2000



To my beloved parents.....

Abstract of thesis presented to the Senate of Universiti Putra Malaysia in
fulfilment of the requirements for the degree of
Master of Agricultural Science

DEVELOPING A DIET OF CATERPILLARS AND MEALWORMS
FOR MASS REARING OF THE PREDATORY BUG
CANTHECONIDEA FURCELLATA (WOLFF.)
(HEMIPTERA: PENTATOMIDAE)

By

EVAWATY SRI ULINA

September 2000

Chairman : Professor Dr. Khoo Khay Chong

Faculty : Agriculture

The asopin bug, *Cantheconidea furcellata* Wolff., is a promising biological control agent against leaf-eating caterpillars of oil palms. Therefore, the bugs should be mass reared efficiently, effectively and economically. A major challenge to the rearing of the predatory bugs is to overcome the difficulty of keeping them fed with live prey. One approach to improve mass rearing is to reduce the dependence on live caterpillars as a food source.

A food item that is readily available and accepted by *C. furcellata* is the mealworms (larvae of *Tenebrio molitor*). In a series of experiments, mealworms were compared with the larvae of *Spodoptera litura* as food

sources. The larvae of *S. litura* were evaluated live, frozen, reared on artificial diet and natural diets. The results showed that larvae of *S. litura* reared on either fresh foliage or artificial diets were the best food sources and that mealworms should be used as a supplementary source.

The second part of the study determined the stage of the nymph during which it is critical to provide live larvae of *S. litura*. The results suggest that developmental and survival rates of second instar nymphs fed live larvae of *S. litura* was comparable to the control (bugs fed live larvae throughout the nymphal stage). However, in terms of the weight of adult females, the control was the best.

The third and final part of the study evaluated the effect of nymphal diet on the reproductive performance of the female bugs. In terms of fecundity and hatchability, feeding the second instar of *C. furcellata* with live larvae of *S. litura* was just as good as feeding the entire nymphal stage with the larvae.

It can be concluded that it is only necessary to feed the second instar nymphs of *C. furcellata* with quality food, i.e. the live larvae of *S. litura*, while other instars may be fed mealworms without affecting growth, survival, fecundity and hatchability of *C. furcellata*.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia
sebagai memenuhi keperluan untuk ijazah Master Sains Pertanian

**MEMAJUKAN SUMBER PEMAKANAN DARI ULAT DAN ULAT
TEPUNG BAGI PEMBIAKAN BESAR-BESARAN KEPINDING
PEMANGSA *CANTHECONIDEA FURCELLATA* (WOLFF.)
(HEMIPTERA: PENTATOMIDAE)**

Oleh

EVAWATY SRI ULINA

September 2000

Pengerusi : Professor Dr. Khoo Khay Chong

Fakulti : Pertanian

Kepinding jenis asopin, *Cantheconidea furcellata* Wolff., merupakan agen kawalan biologi ulat pemakan daun kelapa sawit. Oleh itu, kepinding ini harus dipelihara secara besar-besaran dengan cekap, berkesan dan ekonomik. Namun begitu, kesulitan untuk membekalkan mangsa hidup sebagai makanan kepinding ini merupakan suatu cabaran utama yang perlu diatasi terlebih dahulu. Satu pendekatan untuk memperbaiki pembiakan besar-besaran ialah dengan mengurangkan ketergantungan kepada ulat pemakan daun yang hidup sebagai sumber pemakanan.

Bahan makanan yang tersedia ada dan boleh diterima oleh *C. furcellata* ialah sejenis ulat tepung (larva *Tenebrio molitor*). Dalam beberapa siri kajian yang

dilakukan, ulat tepung telah dibandingkan dengan larva *Spodoptera litura* sebagai sumber pemakanan. Penilaian telah dilakukan ke atas larva *S. litura* yang masih hidup, dibekukan, dan dipelihara menggunakan diet tiruan dan asli. Hasil penilaian menunjukkan larva *S. litura* yang dipelihara sama ada menggunakan daun segar atau makanan tiruan adalah yang terbaik untuk dijadikan sumber pemakanan dan ulat tepung pula digunakan sebagai sumber pemakanan tambahan.

Bahagian kedua kajian ini menentukan peringkat nimfa yang paling kritikal bagi pembekalan larva *S. litura* yang hidup. Keputusan cadang kadar perkembangan dan kemandirian kepinding yang dibekalkan larva hidup *S. litura* diperingkat nimfa instar kedua adalah setanding dengan rawatan kawalan (kepinding yang dibekalkan larva hidup sepanjang peringkat nimfa). Walau bagaimanapun, dari sudut berat dewasa betina, rawatan kawalan adalah yang terberat.

Bahagian ketiga dan terakhir kajian menilai kesan diet nimfa terhadap keupayaan pembiakan kepinding betina. Dari segi fekunditi dan penetasan, pembekalan larva hidup *S. litura* kepada instar kedua *C. furcellata* adalah sama baiknya dengan pembekalan kepada semua peringkat nimfa yang ada.

Kesimpulan dibuat ialah peringkat instar kedua nimfa *C. furcellata* memerlukan pembekalan bahan makanan berkualiti, yakni larva *S. litura*, manakala peringkat instar yang lain memadai dibekalkan ulat tepung tanpa memberi kesan terhadap tumbesaran, kemandirian, fekunditi dan penetasan *C. furcellata*.

ACKNOWLEDGEMENTS

All praise be to Allah SWT., the Almighty, the Most Gracious and Most Merciful, without Whose will no one can achieve anything, and my salawat and salam to His righteous messenger, prophet Muhammad SAW.

I would like to take this opportunity to express my appreciation and gratitude to the chairman of my Supervisory Committee, Prof. Dr. Khoo Khay Chong for his invaluable suggestions, guidance, discussion and patience throughout the project. I am also very grateful to the other members of my supervisory committee, Assoc. Prof. Dr. Yusof Ibrahim and Assoc. Prof. Dr. Dzolkhifli Omar for their constructive comments towards the project and preparation of this thesis.

My sincere gratitude is also extended to the financial support provided by the IRPA fund for this research, which was awarded to Prof. Dr. Khoo Khay Chong. Acknowledgement is also due to all lecturers and staff of Plant Protection Department for their generous co-operation, especially Dr. Mohd. Matthieu Abdullah for his guidance and constructive comments and Mr. R. Selvarajan for his assistance throughout the project. To all my housemates and friends in Indonesian Student Association who had given me the moral encouragement and support to complete my graduate study

and for making my days in Malaysia a memorable and enjoyable one. I would like to extend my thanks to the MARDI, Serdang for their co-operation.

Finally, I am forever indebted to my beloved parents who have given me encouragement and always being immense pillars of support at all times. To my loving brothers for their prayers and support during the many years of my seemingly never ending pursuit of knowledge. I am forever grateful to all my loved ones for being a great source of encouragement from the beginning till the end.

I certify that an Examination Committee met on 11 September, 2000 to conduct the final examination of Evawaty Sri Ulina on her Master of Agriculture Science thesis entitled "Developing a Diet of Caterpillar and Mealworm for Mass Rearing of the Predatory Bug *Cantheconidea furcellata* Wolff. (Hemiptera: Pentatomidae)" in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that candidate be awarded the relevant degree. Members of the Examination Committee are as follows:

ROSLI BIN MOHAMAD, Ph.D.,
Professor,
Faculty of Agriculture,
Universiti Putra Malaysia.
(Chairman)

KHOO KHAY CHONG, Ph.D.,
Professor,
Faculty of Agriculture,
Universiti Putra Malaysia.
(Member)


YUSOF BIN IBRAHIM, Ph.D.,
Associate Professor,
Faculty of Agriculture,
Universiti Putra Malaysia.
(Member)

DZOLKHI FLI BIN OMAR, Ph.D.,
Associate Professor,
Faculty of Agriculture,
Universiti Putra Malaysia.
(Member)


MOHD. GHAZALI MOHAYIDIN, Ph.D.,
Professor/Deputy Dean of Graduate School,
Universiti Putra Malaysia

Date: 10 OCT 2000


This thesis was submitted to the Senate of Universiti Putra Malaysia and was accepted as fulfilment of the requirements for the degree of Master of Agriculture Science.


KAMIS AWANG, Ph.D.,
Associate Professor,
Dean of Graduate School,
Universiti Putra Malaysia.

Date: 11 JAN 2001

DECLARATION

I hereby declare that the thesis is based on my original work except for quotations and citation, which have been duly acknowledge. I also declare that it has not been previously or concurrently submitted for any other degree at Universiti Putra Malaysia or other institutions.



EVAWATY SRI ULINA

Date: 4 OCT 2000

TABLE OF CONTENTS

	Page
DEDICATION	ii
ABSTRACT	iii
ABSTRAK	v
ACKNOWLEDGEMENTS	viii
APPROVAL SHEETS	x
DECLARATION FORM	xii
LIST OF TABLES	xvi
LIST OF FIGURES	xxii
 CHAPTER	
I INTRODUCTION	1
II LITERATURE REVIEW	7
Malaysian Oil Palm	7
Major Oil Palm Pest in Malaysia	9
Integrated Pest Management in Oil Palm	11
<i>Cantheconidea furcellata</i>	13
Distribution	13
Biology	14
Prospect for use in biological control	17
Factors Affecting Insect Growth and Development in Culture	18
Temperature	18
Nutrition	20
Food Consumption in the Immature Stages	23
Body size and weight	23
Stadial duration	26
Food Consumption in the Mature Stage	29
Mating	30
Production of offspring	32
Alternative Sources of Food	36
Factitious host	36
Frozen host	37
Artificial diet	39



	<i>Spodoptera litura</i> Fabr.	41
	<i>Tenebrio molitor</i> L.	42
III	GENERAL MATERIALS AND METHOD	44
	Stock culture of <i>Cantheconidea furcellata</i>	44
	Stock culture of <i>Spodoptera litura</i>	46
	Artificial Diet	47
	Preparing Frozen Larvae of <i>Spodoptera litura</i>	48
	Mealworms	48
IV	EFFECT OF FEEDING PREDACEOUS NYMPHAL INSTARS OF <i>CANTHECONIDEA FURCELLATA</i> WITH LARVAE OF <i>SPODOPTERA LITURA</i> AND <i>TENEBRIO MOLITOR</i>	49
	Introduction	49
	Materials and Methods	50
	Methodology	50
	Statistical analyses	51
	Experiment 1. <i>Cantheconidea furcellata</i> nymphs fed larvae of <i>Spodoptera litura</i> and mealworms	51
	Method	51
	Results	51
	Experiment 2. <i>Cantheconidea furcellata</i> nymphs fed live and frozen larvae of <i>Spodoptera litura</i>	55
	Method	55
	Results	56
	Experiment 3. <i>Cantheconidea furcellata</i> nymphs fed <i>Spodoptera litura</i> reared on fresh foliage and artificial diet	60
	Method	60
	Results	60
	Experiment 4. <i>Cantheconidea furcellata</i> nymphs fed frozen larvae of <i>Spodoptera litura</i> and live mealworms	64
	Method	64
	Results	65
	Discussion	69

V	EFFECT OF FEEDING ADULTS OF <i>CANTHECONIDEA FURCELLATA</i> WITH LARVAE OF <i>SPODOPTERA LITURA</i> AND <i>TENEBRIO MOLITOR</i> ON ITS REPRODUCTIVE PERFORMANCE	73
	Introduction	73
	Materials and Methods	74
	Methodology	74
	Statistical analyses	75
	Results	75
	Discussion	78
VI	NYMPHAL INSTARS OF <i>CANTHECONIDEA FURCELLATA</i> DURING WHICH LARVAE OF <i>SPODOPTERA LITURA</i> IMPORTANT AS FOOD SOURCE	81
	Introduction	81
	Materials and Methods	82
	Methodology	82
	Statistical analyses	84
	Results	84
	Discussion	90
VII	EFFECT OF NYMPHAL FEEDING HISTORY ON REPRODUCTIVE PERFORMANCE OF <i>CANTHECONIDEA FURCELLATA</i>	92
	Introduction	92
	Materials and Methods	93
	Methodology	93
	Statistical analyses	93
	Results	94
	Discussion	97
VIII	GENERAL DISCUSSION	99
	REFERENCES	103
	APPENDICES	121
	BIODATA OF AUTHOR	129

LIST OF TABLES

Table	Page
1 World production of palm oil by major countries (million tonnes)	8
2 Export of Malaysian palm oil to main countries	9
3 Formula for <i>S. litura</i> artificial diet	47
4 Mean developmental times (days \pm SE) of the predaceous nymphs of <i>C. furcellata</i> fed larvae of <i>S. litura</i> and mealworms	52
5 Mean percent mortality (\pm SE) of the predaceous nymphs of <i>C. furcellata</i> fed larvae of <i>S. litura</i> and mealworms	54
6 Mean body weight (mg \pm SE) of adult males and females of <i>C. furcellata</i> after being fed larvae of <i>S. litura</i> and mealworms during the nymphal stage	55
7 Mean developmental times (days \pm SE) of the predaceous nymphs of <i>C. furcellata</i> fed live and frozen larvae of <i>S. litura</i>	57
8 Mean percent mortality (\pm SE) of the predaceous nymphs of <i>C. furcellata</i> fed live and frozen larvae of <i>S. litura</i>	59
9 Mean body weight (mg \pm SE) of adult males and females of <i>C. furcellata</i> after being fed live and frozen larvae of <i>S. litura</i> during the nymphal stage	60
10 Mean developmental times (days \pm SE) of the predaceous nymphs of <i>C. furcellata</i> fed larvae of <i>S. litura</i> reared on either fresh foliage of <i>R. communis</i> or artificial diet	61
11 Mean percent mortality (\pm SE) of the predaceous nymphs of <i>C. furcellata</i> fed larvae of <i>S. litura</i> reared on either fresh foliage of <i>R. communis</i> or artificial diet	63

12	Mean body weight (mg \pm SE) of adult males and females of <i>C. furcellata</i> after being fed larvae of <i>S. litura</i> reared on either fresh foliage of <i>R. communis</i> or artificial diet during the nymphal stage	64
13	Mean developmental times (days \pm SE) of the predaceous nymphs of <i>C. furcellata</i> fed frozen larvae of <i>S. litura</i> and live mealworms	66
14	Mean percent mortality (\pm SE) of the predaceous nymphs of <i>C. furcellata</i> fed frozen larvae of <i>S. litura</i> and live mealworms	67
15	Mean body weight (mg \pm SE) of adult males and females of <i>C. furcellata</i> after being fed frozen larvae of <i>S. litura</i> and live mealworms during the nymphal stage	68
16	Reproductive performance of <i>C. furcellata</i> females fed larvae of <i>T. molitor</i> or <i>S. litura</i> reared on either artificial diet or fresh foliage of <i>R. communis</i>	76
17	Fecundity and hatchability of eggs produced by <i>C. furcellata</i> females of isolated pairs fed larvae of <i>T. molitor</i> or <i>S. litura</i> reared on either artificial diet or fresh foliage of <i>R. communis</i>	77
18	Mean developmental times (days \pm SE) of the predaceous nymphs of <i>C. furcellata</i> fed different prey combinations	85
19	Mean percent survival (% \pm SE) of the predaceous nymphs of <i>C. furcellata</i> fed different prey combinations	87
20	Mean body weight (days \pm SE) of adult males and females of <i>C. furcellata</i> after being fed different prey combinations during the nymphal stage	89
21	Effect of nymphal diet on reproductive performance of <i>C. furcellata</i> females	94
22	Effect of nymphal diet on fecundity and fertility of eggs produced by <i>C. furcellata</i> females	96

LIST OF FIGURES

Figure		Page
1	Adult of <i>C. furcellata</i>	16
2	Diagram illustrating the various pathways by which food intake can influence egg maturation	34
3	Experimental cage for rearing <i>C. furcellata</i>	45
4	Various prey combinations offered to nymphs of <i>C. furcellata</i>	83

CHAPTER I

INTRODUCTION

Oil palm is one of the most important plantation crops in Malaysia. In 1996, the country had more than 2.6 million ha of oil palm and produced 8.39, 1.11, 1.38 and 0.91 million tonnes of palm oil, palm kernel oil, palm kernel cake and oleochemicals, respectively. The exported products earned Malaysia RM 11.51 billion (Porla, 1997). The large export values illustrate the significant contributions of the oil palm industry in the socio-economic development of Malaysia.

The Gross National Product (GNP) could be higher if the oil palm industry can reduce the cost of production. According to Davidson (1991) improved agronomic practices could reduce or maintain the cost of production. An efficient and effective management system for oil palm pests could also reduce cost considerably.

Various pests attacking the oil palm have been described by Wood (1968). These include insects, particularly beetles and caterpillars, some mites and vertebrates. Amongst these pests, the defoliating caterpillars are the

major pests which have caused extensive outbreaks (Teh, 1996). Actually, these pests are classified as occasional pests (Hoong and Hoh, 1992; Chung *et al.*, 1995). They are normally kept under good control by natural enemies, but may flare up from time to time in the absence of natural control. When present in large numbers, they can cause extensive defoliation, eventually affecting yield. Hoong and Hoh (1978) reported that defoliation by nettle caterpillar (*Setora nitens*) caused a reduction in yield of 27 tonnes per ha of fresh fruit bunch (ffb) in severely defoliated palms (above 60% defoliation) during a 30 month period.

There are several factors that influence the increase of a pest population to outbreak levels. The main causes are unhealthy cultural practices and intensive spraying with contact-action insecticides (Hoong and Hoh, 1992; Teh, 1996; Chung, 1998).

Integrated pest management (IPM) is considered to be a suitable approach to avoid outbreaks of leaf-eating caterpillars in oil palm (Chung *et al.*, 1995; Chung, 1998). In IPM, recommendations for pest control are based on an ecological approach. In this way natural enemies are conserved and acute and chronic outbreaks of leaf-eating caterpillars can be avoided.

The term IPM has been defined by many experts. For example, Smith and Reynold (1966) defined IPM as: "A pest management system that in the context of the associated environment and the population dynamics of the pest species, utilises all suitable technique and methods in as compatible a manner as possible and maintains the pest populations at levels below those causing economic injury". The Council on Environment Quality (1972) explained that IPM is an approach that employs a combination of techniques to control the wide variety of potential pests that may threaten crops. It involves maximum reliance on natural pest population controls, along with a combination of techniques that may contribute to suppression - cultural methods, pest-specific diseases, resistant crop varieties, sterile insects, attractants, augmentation of parasites or predators, or chemical pesticides as needed. Flint and van den Bosch (1981) defined IPM as an ecologically based pest control strategy that relies heavily on natural mortality factors such as natural enemies and weather and seeks out control tactics that disrupt these factors as little as possible.

Biological control is an important component of IPM. Biological control can be defined as the action of parasites, predators, or pathogens in maintaining another organism's population density at a lower average than would occur in their absence (De Bach, 1964). In practice it involves (a) the

conservation or maximisation of the role of indigenous organisms in the pest population, (b) repeated release of a biocontrol agent, or (c) single or occasional release of organisms or gene products.

As already explained, leaf-eating caterpillars of oil palm are normally controlled by their natural enemies, including parasitoids, predators and entomopathogens. The common parasitoids are those from the family Chalcidoidea, Ichneumonidae, Braconidae and the order Diptera (Cock, 1987; Desmier de Chenon *et al.*, 1989). Amongst the fungal pathogens infecting leaf-eating caterpillars of oil palm, *Cordyceps* is the most common in Indonesia (Desmier de Chenon *et al.*, 1989) and Malaysia (Mackenzie, 1977). Viruses as entomopathogens belong to several groups, such as Baculoviridae, Reoviridae, Iridoviridae, Parvoviridae and Picornaviridae (Entwistle, 1987). The distribution of these pathogenic viruses are from Indonesia (Kalshoven, 1981; Desmier de Chenon, 1990) to Malaysia (Tiong and Munroe, 1977) and other places in South-east Asia, West Africa and South America (Entwistle, 1987). The important predators are not only the hemipterans from Pentatomidae, Pyrrhocoridae and Reduviidae (Dolling, 1987) but also from the order Coleoptera (Desmier de Chenon *et al.*, 1989).

Among the important predators of leaf-eating caterpillars, *Cantheconidea furcellata* Wolff. (Pentatomidae, Asopinae), also referred to as *Eocanthecona furcellata*, is the most promising. It lays eggs directly on the foliage of the palms, has high fecundity, nymphs and adults live on young and mature palms, and has a short life cycle of about two months (Sipayung *et al.*, 1992).

The use of indigenous biological control agents for the purpose of augmentation or inundation requires that the species can be successfully mass reared. Successful mass rearing requires that the insect species in question can be reared economically without jeopardising quality. Conditions for the nutrition of natural enemy is of special value in the success of insect mass rearing programmes. Inadequate nutrition usually results in great changes in the metabolism (Woodring *et al.*, 1979) and behaviour (Engelmann, 1970; Agarwala and Dixon, 1992) of the insect. These changes have a negative effect on the insects released. Therefore, improvements in a diet for both immature and mature bugs offer the greatest opportunity for improving the quality of mass rearing.

Normally, rearing of the predatory bug is done by feeding with live herbivorous insects. This technique can be difficult and costly if the